



REVIEW: Role of Therapeutic Touch in the management of behavioural and psychological symptoms of dementia.



Ananthavalli Kumarappah (MSc)^{4,5}; Helen Senderovich (MD, MCFP)^{1,2, 3,4, 5}

¹Baycrest, Toronto, Canada; ²Department Family and Community Medicine, ³Division of Palliative Care, ⁴Faculty of Medicine, ⁵University of Toronto, Canada

BACKGROUND

Behavioural and Psychological symptoms of dementia :

- In 2010, there were more than 36 million individuals worldwide living with dementia.
- An estimated 90% of individuals with dementia experience behavioural symptoms (the ways patients respond to their personal, social, or physical environment) over the course of their illness (eg. actions, words and gestures).
- Behavioural symptoms may be harmful to patient, and can be physically and emotionally tolling on family and other caregivers
- Untreated symptoms lead to disease progression, functional decline, repeated falls, longer hospitalization, misuse of medication, and decline in quality of life

METHODS

- To evaluate the current use of Therapeutic Touch (TT) in managing behavioural and psychological symptoms in patients with dementia, we searched PubMed 'Therapeutic Touch'. We limited our inclusion to reviews studies published in the last 10 years (January 2005 to January 2015). We excluded articles in languages other than English and studies for which no outcomes were reported.

STUDIES REVIEWED

Author (Year)	Study Design	Population	Outcomes (instruments)	Authors' Conclusions
Gregory and Verdouw (2005) ³	Exploratory study with pre-test/post-test assessment	Elderly residents (n=121; 24 with dementia)	Changes in physiology, pain level, and behaviour	88% and 92% of patients with dementia experienced physical improvement and positive behavioural changes, respectively. Most commonly observed behavioural changes included reduced vocalizations, reduced restlessness, and improved demeanor.
Woods, Craven, and Whitney (2005) ⁴	Randomized, double-blind, three-group experimental study	57 residents in (age 67-93) with behavioural symptoms of dementia	Agitated Behaviour Response Scale	Reduced manual manipulation/restlessness and vocalization with TT (ANOVA) (F = 3.331, P = .033) and the Kruskal-Wallis test (chi2 = 6.661, P = .036).
Doherty et al. (2006) ⁵	Focus group with semi-structured interviews (staff), pre-test/post-test assessment (residents)	Staff (n=32), 4 residents	Dementia Care Mapping and Positive Response Schedule	Insufficient evidence for use of TT in dementia; improved staff perceptions.
Hawranik, Johnston, and Deatrich (2008) ⁶	Multiple time series, blinded, experimental design	Residents in long-term care with Alzheimer's disease (n=51)	Cohen-Mansfield Agitation Inventory	Reduced nonaggressive behaviours.
Woods, Beck, and Sinha (2009) ⁷	Double-blind experimental interrupted time series	(n=65)	Agitated Behaviour Rating Scale, salivary and urinary cortisol	Decreased restlessness with TT, greater variability of morning cortisol levels (<0.0001).

THERAPEUTIC TOUCH

- TT is an alternative therapy that was described by Dolores Krieger, PhD, RN and Dora Kunz in the 1970s, but is based on ancient healing practices¹.
- In the centering phase, the healer achieves inner calmness and intends to be of service to the patient. This phase can be achieved by quiet meditation or by taking deep breaths to relax the body.
- During the assessment phase, the healer places her/his hands two to four inches over the patient's body to gauge the energy field of the patient. These movements are performed with the palms facing towards the patient, and gradually move from the head of the patient to the feet of the patient. While performing these movements, the healer needs to be aware of sensory cues such as warmth, coolness, static, and tingling.
- During the treatment phase, the healer uses rhythmical and symmetrical movements of the hands to rebalance any disturbed flows of energy. According to practitioners, this method works because healers are able to realign the body's energy field.
- This mechanism of action is controversial due to its speculative nature. Alternatively, Meehan proposes that TT has mostly a placebo and psychological effect².



RESULTS

- Manual manipulation/restlessness and vocalization were the specific behaviours that showed a statistically significant improvement with TT (ANOVA) (F = 3.331, P = .033) and the Kruskal-Wallis test (chi2 = 6.661, P = .036). It may be that TT studies show improvement in agitation and restlessness since they are among the common symptoms associated with dementia. Lack of power can make it difficult to ascertain the value of TT for other less common behavioural and psychological symptoms of dementia. Understanding the specific behaviours that are improved with TT will be important in bringing TT into clinical practice.
- The double-blind RCT showed no significant differences in cortisol levels between the groups⁴. However, there was a statistically significant difference in the morning cortisol variability among the groups (<0.0001). Elevated cortisol levels was observed in dementia. TT was relaxing the patient and reducing their cortisol levels which may slow down disease progression. Further research measuring cortisol levels is necessary to elucidate these results.
- TT practitioners commented on improved therapeutic relationships and ability to maintain a form of communication that may not otherwise be possible in patients with dementia⁵.
- In addition to TT, other forms of touch have been studied as alternative therapies in managing behavioural responses including massage, hand stroking, and holding hands. Studies that further examine TT and other touch modalities are necessary to separate the benefits of TT vs other touch modalities.

CONCLUSIONS

- Despite limited evidence, TT may be explored as an adjunctive therapy for controlling behavioural symptoms in individuals with dementia since it is associated with the absence of negative sequelae.
- TT may be beneficial for common behaviours of agitation such as wandering, restlessness, and vocalizations.
- Future studies should include categorizing individuals by dementia sub-type and severity, having larger sample sizes, and better methodological consistency in recording behavioural symptoms.
- For inquiry: please contact Dr. Helen Senderovich at hsenderovich@baycrest.org

REFERENCES

- Krieger, D. *Accepting Your Power to Heal: The Personal Practice of Therapeutic Touch*. (Bear & Company, 1993).
- Meehan, T. C. Therapeutic touch as a nursing intervention. *J. Adv. Nurs.* **28**, 117-125 (1998).
- Gregory, S. & Verdouw, J. Therapeutic touch: its application for residents in aged care. *Aust. Nurs. J.* **12**, 23-5 (2005).
- Woods, D. L., Craven, R. F. & Whitney, J. The effect of therapeutic touch on behavioural symptoms of persons with dementia. *Altern. Ther. Health Med.* **11**, 66-74 (2005).
- Doherty, D., Wright, S., Aveyard, B. & Sykes, M. Therapeutic touch and dementia care: an ongoing journey. *Nurs. Older People* **18**, 27-30 (2006).
- Hawranik, P., Johnston, P. & Deatrich, J. Therapeutic touch and agitation in individuals with Alzheimer's disease. *West. J. Nurs. Res.* **30**, 417-434 (2008).
- Woods, D. L., Beck, C. & Sinha, K. The effect of therapeutic touch on behavioural symptoms and cortisol in persons with dementia. *Forsch. Komplementmed.* **16**, 181-189 (2009).